

RESTRICTED USE PESTICIDE

DUE TO HAZARD TO NON-TARGET SPECIES

For retail sale to and use only by Certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Prozap[®] Zinc Phosphide Oat Bait

EPA Reg. No. 61282-14

SLN No. WY-020003

Active Ingredient: Zinc Phosphide	2.0%
Other Ingredients.....	98.0%
Total.....	100.0%

SPECIAL LOCAL NEED SUPPLEMENTAL LABEL

For Distribution and Use Only in the State of Wyoming

For Control of Prairie Dogs

in Rangeland and Adjacent Timber

Only for Uses and Application Methods on this Supplemental Label

This label valid until April 30th, 2017 or until otherwise amended, withdrawn, cancelled, or suspended.

This label, the container label, and the booklet label for the Federally registered product, Prozap[®] Zinc Phosphide Oat Bait (EPA Reg. No. 61282-14), must be in the possession of the user at the time of application. All applicable directions, restrictions, and precautions on the container label and booklet for the Federally registered product and on this supplemental label must be followed.

DIRECTIONS FOR USE

USE RESTRICTIONS: This product may be used to control black-tailed prairie dogs (*Cynomys ludovicianus*), white-tailed prairie dogs (*C. leucurus*), Columbian ground squirrels (*Spermophilus columbianus*), and Richardson's ground squirrels (*S. richardsonii*) in rangelands and adjacent timber areas in Wyoming only. This product may be used to control prairie dogs from March 1 through December 31.

Before baiting for prairie dogs, contact the U.S. Fish and Wildlife Service office in Cheyenne, WY, at 307-772-2374 to determine whether a black-footed ferret survey is required.

Apply toxic bait no more than once per season. Apply toxic bait only at times of year when target species are accepting grain readily.

Do not apply this product to croplands. Do not apply this product on roads, near residential or public use areas, or over water. Do not apply this bait at any location where endangered species occur or where consumption of prebait by nontarget wildlife was observed.

PREBAITING: Prebait with 4 grams (one teaspoonful) of untreated oat groats per mound or burrow opening, one to two days prior to the anticipated time of baiting. Prebaiting typically increases acceptance of treated baits by prairie dogs and ground squirrels. Observe prebaited areas for evidence of consumption of prebait by target and nontarget species.

BAITING: Apply toxic bait only after all or most of prebait has been eaten and only in areas where prebait was consumed. Apply bait by hand (wear protective gloves) as a six-inch bait spot located on the edges of each mound or burrow opening, or in adjacent feeding areas. Do not apply more than one teaspoonful of bait (approximately 4 grams) per bait spot. Dispose of spilled or excess bait by burial.

24(c) Registrant:
HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956