

RESTRICTED USE PESTICIDE

DUE TO HIGH ACUTE INHALATION TOXICITY AND EYE AND SKIN CORROSIVENESS TO HUMANS; HIGH ACUTE TOXICITY TO NONTARGET BIRDS AND AQUATIC INVERTEBRATES AND THE NEED FOR HIGHLY SPECIALIZED APPLICATOR TRAINING.

For sale and transfer to, and use only by, certified applicators who are USDA APHIS personnel trained in bird control or by persons under their direct supervision.

SUPPLEMENTAL LABELING

COMPOUND DRC-1339 CONCENTRATE— Livestock, Nest & Fodder Depredations -Wyoming EPA Registration Number 56228-29

SLN WY-11-0002

FOR DISTRIBUTION AND USE ONLY WITHIN WYOMING

For control of ravens, crows and magpies that cause threats to human health and safety, livestock, sensitive wildlife species, or structure damage.

THIS LABEL AND THE LABEL FOR THE FEDERALLY REGISTERED PRODUCT, EPA Reg. No. 56228-29, MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. READ THIS ENTIRE LABEL AND THE ENTIRE LABEL FOR 56228-29. FOLLOW ALL DIRECTIONS, RESTRICTIONS AND PRECAUTIONS FOUND ON THIS LABEL AND ALL APPLICABLE REQUIREMENTS ON THE LABEL FOR 56228-29.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and the label for EPA Registration Number 56228-29. This product must be used strictly in accordance with the precautionary statements and applicable use directions on both labels, as well as with all applicable State and Federal laws and regulations.

Before using this product, contact the U.S. Fish and Wildlife Service and the applicable State wildlife agency to obtain all necessary kill or collecting permits.

GENERAL INFORMATION: This product contains a slow-acting avicide which kills target birds in 1 to 3 days. As many types of nontarget bird species are potentially vulnerable to DRC-1339, it is necessary to use care and to follow the requirements of this label to minimize impacts to nontarget species.

USE RESTRICTIONS: Baits made from Compound DRC-1339 Concentrate-Livestock, Nest & Fodder Depredations-Wyoming may only be used to control:

Common ravens (*Corvus corax*)
American crows (*Corvus brachyrhynchos*)
Black-billed magpies (*Pica pica*)

This product may be used only to prepare dry dog food and dry cat food baits to control species listed on this label.

–**DO NOT** apply baits made from this product by air or by using any procedures not prescribed on this label.

–This product may only be used to make baits with dry dog food or dry cat food baits. Prepare baits following the instructions on this label under **BAIT PREPARATION AND APPLICATION**.

–**DO NOT** use treated baits as food or feed.

–**DO NOT** apply this product in any way that could contaminate food, feed or water, or that would allow bait to be consumed by livestock.

–**DO NOT** graze livestock or plant any crop in treated noncrop areas or in crop areas that may have been contaminated by treated bait for one year (365 days) following the last application of bait made from this product.

– **DO NOT exceed a maximum single application rate of 0.1 lb of active ingredient per treated acre or a maximum cumulative application rate of 0.5 lb of active ingredient per treated acre in any use year.** The maximum single treatment rate of 0.1 lb of active ingredient per acre would be provided by one treatment with a 1.9% DRC-1339 bait diluted 1:10 with untreated bait material and applied at a rate of 58 lbs of diluted bait per acre.

–**DO NOT** apply treated baits within 50 feet of bodies of water.

–Before baits made from this product are applied, sites that are to be treated must be observed for evidence of nontarget activity and must be prebaited (see specific instructions for these activities). **DO NOT** apply treated baits in locations where prebait has not been accepted well by target species or where nontarget wildlife have been observed to feed on prebait.

–**DO NOT** apply bait(s) in areas where there is danger that Threatened or Endangered species will consume baits, unless special precautions are taken to limit such exposures. Such precautions shall include constant observation of baited sites and hazing techniques to frighten away Threatened and Endangered species that otherwise might feed on baits. Note: A permit may be required for harassment of Threatened or Endangered Species. See **THREATENED AND ENDANGERED SPECIES** section on the Federal label for 56228-29 for information on consideration of threatened and endangered species.

–**DO NOT** use bait materials that resemble human foods in any areas that are accessible to children.

–**DO NOT** apply baits made from this product in a way that will contact workers or other persons.

ENTRY RESTRICTIONS: Only protected applicators may be in the area during bait application. Keep pets and livestock, and persons other than authorized handlers away from the bait at all times.

PRETREATMENT OBSERVATIONS: Prior to application of treated bait, observe habits of target birds to determine their relative numbers, feeding habits, preferred feeding locations, daily activity patterns, and the relative abundance and types of nontarget species at potential baiting sites. Once potential baiting sites are identified, they must be observed carefully to determine the potential hazards to nontarget species and must be prebaited to promote bait acceptance by target species. **DO NOT** apply treated bait where there is a danger that it will be consumed by Threatened or Endangered species.

PREBAITING: Prebaiting with untreated bait material of the type to be used for toxic baiting is required. After daily activity patterns of target species have been noted and the appropriate baiting location has been selected, apply prebait using the same application method and application rate that is to be used for toxic baiting (see **BAIT PREPARATION AND APPLICATION DIRECTIONS**).

BAIT PREPARATION AND APPLICATION DIRECTIONS:

Dry dog or dry cat food baits:

- Place 10 lbs (4.5 kg) of dry dog food nuggets or dry cat food into a mixer, plastic container or a leak-proof plastic bag and set aside.
- Warm a mixture of 333 ml (10 oz) of edible oil and 80 ml (3 oz) of Alcolec-S (lecithin) to 110 °F (43 °C) for about 5 minutes. Dissolve 46 g (1.6 oz) of Compound DRC-1339 Concentrate - Staging Areas and Feedlots - WY in the oil mixture.
- Pour resulting solution or suspension over the 10 lbs (4.5 kg) of dry dog food nuggets or dry cat food and mix or tumble slowly until the bait is evenly mixed.
- Air dry bait at ambient temperature. When bait is dry, mix treated dry dog food or dry cat food with untreated dry dog food or dry cat food of the same type at a ratio of 1 part of treated nuggets to 10 parts of untreated dry dog food or dry cat food.

POST THE AREA WITH WARNING SIGNS DURING THE BAITING OPERATION IF BAIT IS LEFT UNATTENDED OR IF CHILDREN, PETS OR DOMESTIC ANIMALS HAVE ACCESS TO THE BAITING SITE.

BAIT STABILITY: Treated baits prepared as described above and stored under cool and dry conditions may be used up to 30 days after preparation. Use diluted baits within 7 days after preparation. Immediately use treated or diluted baits that have been exposed to sunlight or heat in excess of 110° F (43°C) following preparation, or dispose of them as pesticide waste.

APPLICATION DIRECTIONS:

Wear rubber gloves while handling treated baits and bait mixtures and dispense or broadcast baits using a scoop or other appropriate equipment.

Application Rates: Do not make any treatment which exceeds rates of 58 lbs/treated acre dry dog food or dry cat food baits. Do not make more than 5 treatments per year to any one treated site.

MANUAL APPLICATIONS (NON-CROP AREAS ONLY):

-When target birds are observed feeding well on the prebait and about 75% of the prebait has been consumed, apply treated baits.

-Baits may be distributed on the ground, or placed in bait stations, on feeding trays or on elevated platforms. DO NOT exceed a maximum application rate of 0.1 lb (1.6 oz) of active ingredient per acre or a cumulative treatment rate of 0.5 lb (8 oz) of active ingredient per acre for the use year. A single application rate of 0.1 lb (1.6 oz) of active ingredient per acre is consistent with use of 1) 110 lbs (49.9 kg) of diluted dry dog food or dry cat food. To determine the cumulative treatment rate of 0.5 lb (8 oz) of active ingredient per acre per use year, multiply the maximum amount of bait that can be applied per acre per treatment by 5.

POST-TREATMENT CLEAN-UP: Collect and remove all unconsumed bait from bait stations at the conclusion of the treatment period. Dispose of recovered and unused bait in accordance with the label for 56228-29 and applicable Federal, State, and local laws. Beginning one day after bait application, search treated areas, immediate surrounding areas, and known roost sites (if accessible). Collect carcasses of dead or dying birds and bury or burn them, according to applicable laws. Repeat carcass searches at 1-2 day intervals as long as bait is exposed and likely to remain toxic.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store only in original container, in a dry place inaccessible to children, pets, and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bags by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. If bags are not to be recycled, dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities. If burned, stay out of smoke.

EPA SLN NUMBER WY-11-0002

Label Issue Date:

Label Expiration Date:

24(c) Registrant:

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
4700 River Road, Unit 149
Riverdale, MD 20737-1237

Local Contact:

State Director
USDA APHIS
Wildlife Services
6731 West Coal Road
Casper, WY 82604

Rev: 03/16/2011